



**PRODUCT INFO
& DATASHEET**

SICOHEL-DELAY

*Chelated Micronutrients Mix and Plant Growth Regulator (PGR)
Water-soluble powder for foliar and soil application*

1. DESCRIPTION

SICOHEL-DELAY contains the essential micro-nutrients boron (B), copper (Cu), iron (Fe), manganese (Mn), molybdenum (Mo) and zinc (Zn) as well as coadjuvants for setting and fruiting.

SICOHEL-DELAY is fully water soluble and it can be applied via foliage and soil.

This product is 100% chelated: Chelating agents Na4 EDTA and Citric Acid.

2. GUARANTEED ANALYSIS

		<u>Formula nr. 1</u>		<u>Formula nr. 2</u>	
Boron	(B)	5 g/kg	0.5 %	5 g/kg	0.5 %
Copper	(Cu)	5 g/kg	0.5 %	5 g/kg	0.5 %
Iron	(Fe)	100 g/kg	10.0 %	40 g/kg	4.0 %
Manganese	(Mn)	40 g/kg	4.0 %	40 g/kg	4.0 %
Molybdenum	(Mo)	5 g/kg	0.5 %	5 g/kg	0.5 %
Zinc	(Zn)	10 g/kg	1.0 %	80 g/kg	8.0 %
Contains also:	Naphthyl Acetic Acid (NAA)		0.18 %		
	Naphthylacetamide (NAD)		0.48 %		

3. SUGGESTED APPLICATIONS AND RATES OF USE

* **HORTICULTURE:** aubergines, melons, watermelons, tomatoes, marrows, etc.

Apply every 15-20 days to achieve good setting and fruiting.

Rate: max. 4-5 kg per hectare via fertigation: max. 1-1.5 g/ltr for foliar application.

* **FRUIT:** to aid setting of peaches, apples, pears, stone and pip fruits in general.

Apply every 10-15 days before start of flowering.

Rate: 4-5 kg per hectare repeated every 10-15 days until petal fall.

* **WHEN TO SPRAY:** to get the best results from your feeding programme, do not apply the product during the hottest part of the day. Apply foliar feeds early in the morning or late afternoon.

4. COMPATIBILITY

SICOHEL-DELAY is generally compatible with the majority of chemical crop protection products.

It can be incorporated into existing spray programmes. Add the micronutrients to the water before adding any other ingredients.

It is always advisable to carry out a small scale compatibility trial before mixing a full batch.

5. PACKING

10 kg bags. Other packings on request.