



**PRODUCT INFO  
& DATASHEET**

# SICOGREEN<sup>®</sup>-L SUPER

## Liquid NPK 10-10-7.5 + TE (w/v)

*SICO's well-balanced N-P-K liquid fertiliser for high value crops  
For horticulture, agriculture and hydroponics*

02/2016

### 1. PRODUCT DESCRIPTION

**SICOGREEN<sup>®</sup>-L SUPER** is a liquid fertilizer with a full complement of macro and micronutrients. It is used as a general supplementary foliar feed in horticulture and agriculture for the prevention or elimination of nutrient deficiencies in a wide range of crops. **SICOGREEN<sup>®</sup>-L SUPER can also be used as soil fertiliser (esp. in horticulture), and has proved to be very successful for hydroponics.** Due to the additives in the formulation, the rate of nutrient uptake is high. As a consequence, it is particularly suitable for improving the leaf colour and shine of ornamentals, even shortly before they are sold. And it does not leave marks!

### 2. CONTENTS

Macronutrients		% w/v **	% w/w
Total N (nitrogen)		10	8
Nitrate N	NO <sub>3</sub> -N	2.8	2.3
Ammoniacal N	NH <sub>4</sub> -N	4.6	3.7
Carbamide N	NH <sub>2</sub> -N	2.5	2.0
P <sub>2</sub> O <sub>5</sub> (phosphate)	water soluble	10	8
K <sub>2</sub> O (potassium)	water soluble	7.5	6
Micronutrients		% w/v	% w/w
B (boron)		0.124	0.01
Cu* (copper)		0.0087	0.007
Fe* (iron)		0.0187	0.015
Mn* (manganese)		0.0161	0.013
Mo (molybdenum)		0.0012	0.001
Zn* (zinc)		0.0062	0.005
		% w/v	% w/w
Cl <sup>-</sup>		1.1	0.9
Na <sup>+</sup>		< 0.1	< 0.1

\* Fully chelated micronutrient (EDTA)

\*\* Calculated on density

### 3. PHYSICOCHEMICAL PROPERTIES

Appearance:	green liquid			
Density (20°C/68°F):	approx. 1.24			
pH (20°C/68°F):	approx. 5.4			
Conductivity in distilled water (25°C/77°F):				
°/°°	0.5	1.0	1.5	2.0
µS/cm	270	510	770	990

### 4. PRECAUTIONS

When storing **SICOGREEN<sup>®</sup>-L SUPER**, temperatures below +5°C (41°F) and above +40°C (104°F) should be avoided. Too low temperatures (and long-term storage) can cause crystallization. The crystals will however easily dissolve again in the spray solution. When mixing with pesticides for the first time, test on a small scale before general use.

### 5. BIOLOGICAL/TECHNICAL PROPERTIES

- \* High and well balanced macronutrient supply.
- \* Complete micronutrient supply.
- \* Suitable for all crops in all climatic zones.
- \* **pH regulation of the spray solution.**
- \* Nutrients readily available to plants.
- \* Very safe to plants.

Any information in this publication is believed to be accurate and is given in good faith, but is for the customer to satisfy himself of the suitability for his own particular purpose. No representation, warranty or guarantee is made to its accuracy, reliability or completeness.

**SAP INTERNATIONAL CORPORATION bvba Krekelenberg 69, B-2980 Zoersel, Belgium**  
Tel. +32-3-309.06.51 Fax. +32-3-309.19.31 Email : [info@sico.be](mailto:info@sico.be) Website : [www.sico.be](http://www.sico.be)

**SICO FERTILISERS**  
EVERY TIME THE *right* SOLUTION



- \* Fully chelated cationic micronutrients.
- \* Superchelation reduces the water hardness of the spray solution.
- \* The optimal foliar fertiliser for all ornamentals: Leaves no leaf spots, improves the leaf colour, achieves good leaf finish, and ensures seedling production.
- \* Improves growth, flowering and fruit performance.
- \* Compatible with most commonly used pesticides.
- \* Can be applied with all usual HV and LV spraying and sprinkling equipment.
- \* **Maintains healthy and attractive leaves.**

## **6. RATES AND METHODS OF USE**

\* We have also indicate rates in ml/15 ltr knapsack sprayer

Crop	Rate	* ml/15 ltr	Number and timing of applications
Agriculture	5-10 l/ha	37.5-75	General recommendation for foliar nutrition in conjunction with all pesticide treatments
Fruit growing	0.3%	45	General recommendation for foliar nutrition and post-blossom spray in conjunction with pesticide treatments approx. 6-7 sprays
Grapes	0.3%	45	For foliar nutrition in conjunction with all pesticide treatments
Potted plants/ Cut flowers	0.05-0.5%	7.5-75	General recommendation for soil and foliar nutrition, either alone or in conjunction with pesticide treatments: watering, sprinkling, spraying or atomizing
* Young plants	0.05-0.1% 0.05%	7.5-75 7.5	Water or spray once to twice a week After true leaves have formed or with each atomizer/spray application
* Adult plants	0.2%	30	Use alone or in conjunction with pesticide treatments; spraying or sprinkling
* Marketable plants	0.5% 0.4%	50 60	As a leaf shine Atomizing
Hydroponics			
* Young plants	0.05%	7.5	Nutrient solution for low nutrient requirements
* Others	0.1-0.2%	15-30	As required
Tree nurseries			
* Multiplication/ nursery stock	0.1-0.2% 0.05%	15-30 7.5	After true foliage leaves have formed spraying or sprinkling or with each atomizer/spray application
* Young plants/ saleable stock	0.2% 0.4%	30 60	Foliar nutrition either alone or in conjunction with pesticide treatments Spraying/sprinkling Atomizing
Vegetables	0.1-0.2% 0.4%	15-30 60	Alone or in conjunction with pesticide treatments, as soil and foliar nutrition. Watering, sprinkling or spraying, + applications at 14 days interval Atomizing, + applications at 14 days interval
Aubergines Cucumbers, Peppers	5 l/ha	37.5	6 applications 1st and 2nd: before flowering 3rd-6th: after flowering Intervals between applications should be 2 weeks throughout
Avocados	0.1% 0.2% 0.4%	15 30 60	6 applications. 2 before flowering 4 after flowering at 2-week intervals Young plants ter planting out Bearing trees
Bananas	4-5 l/ha	37.5	At least 5 applications
Beans, Peas	5 l/ha	37.5	4 applications. 1st: before flowering 2nd-4th: after flowering at 14-day intervals
Brassicas, Lettuce	0.1-0.2%	15-30	4 applications. 1st: after planting out 2nd-4th: at 10-days interval
Carrots	5 l/ha	37.5	6 applications at 14-day intervals
Cereals	5-10 l/ha	37.5-75	In conjunction with pesticide treatments
Citrus	5 l/ha	37.5	6 applications. 1st: in winter 2nd: before the blossoms open - 3rd: after petal fall - 4th-6th: at 4-week interval

Any information in this publication is believed to be accurate and is given in good faith, but is for the customer to satisfy himself of the suitability for his own particular purpose.  
No representation, warranty or guarantee is made to its accuracy, reliability or completeness.



Crop	Rate	ml/15 ltr	Number and timing of applications
Coffee (in plantations)	0.1% 0.2 % 0.4 %	15 30 60	6 applications. 1st: before flowering 2nd-6th: after flowering at 3-week intervals Young plants After planting out Bearing trees
Cotton	5 l/ha	37.5	5 applications. 1st and 2nd: in conjunction with the 1st and 2nd insecticide sprays 3rd-5th: during boll formation at 14-day intervals
Hops	6-8 l/ha	45-60	Before and after flowering, in conjunction with all pesticide treatments
Lucerne, Pasture land Maize	5 l/ha 5-10 l/ha	37.5 37.5	1 application 10 days after each cut 4 applications. 1st: when the plants are 15-25 cm high, 2nd-4th: at 14-days intervals
Manioc	5 l/ha	37.5	4 applications. 1st: 30 days after germination 2nd-4th: at 14-day intervals
Melons, Vegetable Marrows	5 l/ha	37.5	4 applications. 1st: 30 days after germination 2nd-4th: at 14-day intervals
Olives	5-10 l/ha	37.5-75	3 applications. 1st: in early spring 2nd: May/June, 3rd: September/October
Onions	5 l/ha	37.5	4 applications. 1st: 30 days after sprouting 2nd-4th: at 14-day intervals
Papayas	0.2% 0.3%	30 45	6 applications. 1st and 2nd: before flowering at 3-week intervals 3rd-6th: after flowering at 3-week intervals Young plants Bearing trees
Pineapples	5 l/ha	37.5	5 applications. 1st: shortly before blossom 2nd-5th: post-blossom at 20-day intervals
Rice	0.4% 4-5 l/ha	60 37.5	In the seed bed before planting out at least 2 applications Under field conditions, at least 5 applications
Soybeans	5 l/ha	37.5	Several applications 1st: after first flowering, then every 14 days, however 3 applications at least
Strawberries	0.1%- 0.2%	15-30	Weekly applications. 1st: after planting out
Tea	5 l/ha	37.5	6 applications. 1st: when the leaf buds form 2nd-6th: at 3-week intervals
Tobacco	5 l/ha	37.5	Several applications. 1st: at the 4- to 5- leaf stage, 2nd: 3 weeks after the 1st application 3rd: 3 weeks after the 2nd application, etc.
Sugar cane	8 l/ha	60	1-2 applications, at a growth height of 25-35 cm
Sugarbeet	5-10 l/ha	37.5-75	2-4 applications between 2-leaf stage and before crop cover (before the rows close for tractor passage)
Cocoa	5 l/ha (= 0.5%)	37.5	2 - 3 applications
Palm trees (container nurseries)	5 l/ha (= 0.5%)	37.5	Monthly spray
Pistachio			foliar: dilute 200-300 ml/hl of water, soil application: dilute 400 ml/hl of water Use up to 1000 l of water/ha. Make sure plants are well wetted.

## 7. PACKAGING

1 l / 5 l / 10 l / 20 l / 200

Any information in this publication is believed to be accurate and is given in good faith, but is for the customer to satisfy himself of the suitability for his own particular purpose.  
No representation, warranty or guarantee is made to its accuracy, reliability or completeness.

**SAP INTERNATIONAL CORPORATION bvba** Krekelenberg 69, B-2980 Zoersel, Belgium  
Tel. +32-3-309.06.51 Fax. +32-3-309.19.31 Email : [info@sico.be](mailto:info@sico.be) Website : [www.sico.be](http://www.sico.be)