



**PRODUCT INFO
& DATASHEET**

FURAZOLIDONE

ANTIBIOTIC & GROWTH PROMOTOR

DESCRIPTION

Furazolidone waterdispersible 96 % has been designed for convenient administration in the drinking water of the animal. It is especially suitable for animals too sick to eat or for those with a decreased feed intake. Furazolidone is indicated for the treatment of Salmonella infections, especially *S. pullorum* and *S. gallinarum*, and in *E. coli* infections, such as Colisepticaemia. It is also effective in the treatment and prevention of diseases caused by furazolidonesensitive organisms. It is furthermore effective in the treatment and prevention of blackhead (histomoniasis), hexamitiasis and bacterial enteritis. Furazolidone is also useful in cases of sinusitis in turkeys and of great value in chronic respiratory disease (CRD), especially against the secondary infections associated with the disease. Furazolidone is also used for the treatment and prevention of coccidiosis, either alone or in combination with other nitrofurans.

SPECIFICATIONS

Furazolidone, micronized 96 % - 5 micron: 96 %

Suspending agents: 4 %

RECOMMENDED DOSAGES

***Poultry**

Treatment : 1-2 grams of furazolidone, waterdispersible 96 % per 10 liters of drinking water for 3-7 days.

Prevention : 0.5 -1 gram of furazolidone waterdispersible 96 % per 10 liters of drinking water.

Note :

- Only furazolidone-medicated water should be administered during treatment and prevention periods.
- It is important to mix the suspension every few hours, but at least every 8 hours.

***Calves, Sheep and Other animals**

Treatment : 75-100 mg furazolidone, waterdispersible 96 % per 10 kg bodyweight twice daily for 3-4 days.

Prevention : 50-75 mg furazolidone, waterdispersible 96 % per 10 kg bodyweight twice daily.

PACKING

25 kg bags or drums

STORAGE AND HANDLING

Should be stored in a safety place.

CAUTION

- Not to be used in combination with D.O.T. coccidiostat
- Do not administer to waterfowl