



**PRODUCT INFO
& DATASHEET**

SICO® MONENSIN

10 & 20% Anticoccidial & Growth Promotor

INDICATIONS

Prophylaxis of coccidiosis of poultry caused by *Eimeria tenella*, *E. acervulina*, *E. mivati*, *E. brunetti*, *E. maxima*, *E. necatrix*. For fattening of calves and lambs.

CONTRA-INDICATIONS

Fodder, intended for chickens, should not be used for layer hens, turkeys, guinea fowls and other animals, particularly for horses and donkeys. It is incompatible with Tiamulin, Erythromycin, Chloramphenicol, Oleandomycin, Sulfadimidine, Sulfadimethoxine, Sulfaquinoxaline.

MODE OF APPLICATION

Orally, very well mixed with the fodder.

NOTE: for realization of full homogenization of the preparation with the fodder, please follow this recommended scheme for mixing by the method of the stage dilution:

Stage 1: 1 kg of Monensin 10% + 10 kg of fodder

Stage 2: The mixture of stage 1 + 100 kg of fodder

Stage 3: The mixture of stage 2 + 890 kg of fodder

The recommended mixing scheme is given for dosage of Monensin 10% in fowls.

The stage dilution is analogically performed for the other animal species with Monensin 10% and 20% respectively.

DOSAGE

In grams per ton fodder.

Animal	Monensin versus pure substance	Monensin 10%	Monensin 20%
<i>Poultry</i>	90 – 110 g/ton	900 – 1,100 g/ton	450 – 550 g/ton
<i>Calves</i>	5 – 30 g/ton	50 – 300 g/ton	25 – 150 g/ton
<i>Lambs</i>	5 – 10 g/ton	50 – 100 g/ton	25 – 50 g/ton

ADVERSE EFFECTS

Overdosage could inhibit growth and an intoxication is possible, manifested with loss of appetite, tremor, paralysis of the limbs and death.

WITHDRAWAL TIME

3 days.

SHELF LIFE

Two years.

STORAGE

Dry, dark & at room temperature.

PACKING

20 kg bags.